

Ordering Physician:


Metamatrix

3425 Corporate Way

Duluth, GA 30096

2100 Gastrointestinal Function Profile

Methodology: DNA Analysis, GC/MS, Microscopic, Colorimetric, Automated Chemistry, ELISA




Consistency = Formed/Normal


Predominant Bacteria (E+007)

E+007


Obligate anaerobes



Facultative anaerobes



Obligate aerobes



Opportunistic Bacteria

No clinically significant amounts.

Units and Reference Ranges

Organisms are detected by DNA analysis. One colony forming unit (CFU) is equivalent to one bacterium. Each genome detected represents one cell, or one CFU. Results are expressed in scientific notation, so an organism reported as 2.5 E7 CFU/gram is read as 25 million colony forming units per gram of feces. The cutoff for significance of Opportunistic Bacteria has been set at 1.0E+ 005 (100,000). These are levels above which clinically significant growth may be present. Rather than reporting semi-quantitative +1 to +4 levels, the new methodology provides full quantitative analysis.

Predominant Bacteria play major roles in health. They provide colonization resistance against potentially pathogenic organisms, aid in digestion and absorption, produce vitamins and SCFA's, and stimulate the GI immune system. DNA probes allow detection of multiple species (sp.) within a genus, so the genera that are reported cover many species.

Opportunistic Bacteria may cause symptoms and be associated with disease. They can affect digestion and absorption, nutrient production, pH and immune state. Antibiotic sensitivity tests will be performed on all opportunistic bacteria found, although clinical history is usually considered to determine treatment since the organisms are not generally considered to be pathogens.

Ordering Physician:
Metamatrix

3425 Corporate Way
Duluth, GA 30096

Accession Number: **A1106290325**

Reference Number:

Patient: **Sample Report**

Age: 49 **Sex: Female**

Date of Birth: 02/05/1962

Date Collected: 6/27/11

Date Received: 6/29/11

Report Date: 6/29/11

Telephone: (770) 446-4583

Fax: (770) 441-2237

Reprinted: 7/19/11

Comment:

2100 Gastrointestinal Function Profile

Methodology: DNA Analysis, GC/MS, Microscopic, Colorimetric, Automated Chemistry, ELISA

Pathogenic Bacteria 95% Reference Range

Organism	Result	Reference Range
Helicobacter pylori	<0.01	<=1.0E+005
E.H.E. coli	<0.01	<=1.0E+005
Clostridium difficile	<0.01	<=1.0E+005
Campylobacter sp.	<0.01	<=1.0E+005

Yeast/Fungi Expected Value

Candida sp. **+1 => 100 pg DNA/g specimen** Neg

Yeast/Fungi

Yeast overgrowth has been linked to many chronic conditions, in part because of antigenic responses in some patients to even low rates of yeast growth. Potential symptoms include diarrhea, headache, bloating, atopic dermatitis and fatigue. Positives are reported as +1, +2, +3 or +4 indicating >100, >1000, >10000 or >100000 pg DNA/g.

Parasites Expected Value

Blastocystis hominis **Positive** Neg

Parasites

Parasite infections are a major cause of non-viral diarrhea. Symptoms may include constipation, gas, bloating, increased allergy response, colitis, nausea and distention.

Adiposity Index

Phylum	Value	Reference Range
Firmicutes	76	<= 80
Bacteroidetes	24	>= 20

The **Adiposity Index** is derived by using DNA probes that detect multiple genera of the phyla Firmicutes and Bacteroidetes. Abnormalities of these phyla may be associated with increased caloric extraction from food.

Drug Resistance Genes

aacA, aphD	Pos	gyrB, ParE	Neg
mecA	Neg	PBP1a, 2B	Neg
vanA, B, and C	Neg		

Ordering Physician:

Metamatrix


3425 Corporate Way

Duluth, GA 30096


2100 Gastrointestinal Function Profile

Methodology: DNA Analysis, GC/MS, Microscopic, Colorimetric, Automated Chemistry, ELISA

Percentile Ranking by Quintile



Beneficial SCFA



Beneficial SCFA

Short chain fatty acids (SCFA) are produced by bacterial fermentation of dietary polysaccharides and fiber. The product, N-butyrate, is taken up and used to sustain the normal activity of colonic epithelial cells. Butyrate has been shown to lower the risk of colitis and colorectal cancer. A healthy balance of GI microbes depends on production of SCFA by one specie to allow the normal growth of another one in a complex cross-feeding network.


Inflammation



Inflammation

Lactoferrin, an iron-binding glycoprotein, is released in IBD but not in non-inflammatory IBS. High levels are found in Crohn's, UC or infection. WBC's are elevated in general inflammation/infection. Mucus is often visualized in acute GI inflammation.

Immunology



Immunology

High fecal sIgA indicates immune system reactions to the presence of antigens from bacteria, yeast or other microbes. Low sIgA can result from stress or malnutrition. Anti-gliadin sIgA is a screening marker for gluten sensitivity.


Ordering Physician:
 Metamatrix

3425 Corporate Way
 Duluth, GA 30096


2100 Gastrointestinal Function Profile

Methodology: DNA Analysis, GC/MS, Microscopic, Colorimetric, Automated Chemistry, ELISA

Percentile Ranking by Quintile




Additional Tests



Additional Tests

pH is influenced by numerous factors, but it is strongly related to the bacterial release of pH-lowering organic acids and pH-raising ammonia. Positive **RBCs** can signify GI tract bleeding. **Color** (other than brown) abnormalities can be due to upper GI bleeding, or bile duct blockage, steatorrhea or antibiotic use.


Digestion



Digestion

Pancreatic elastase 1 levels below the reference limits are strongly correlated with pancreatic insufficiency. High triglycerides signify fat maldigestion. Putrefactive SCFA are a result of bacterial fermentation of undigested protein. High numbers of vegetable fibers indicate maldigestion.

Absorption



Absorption

High **LCFA** indicates fat malabsorption due to pancreatic or biliary insufficiency, or acute bacterial infection that produces intestinal cell destruction. High total fat usually signals malabsorption, as does elevated fecal cholesterol.

UC** = Unable to Calculate

These test results are not for the diagnosis of disease. They are intended to provide nutritional guidelines to qualified healthcare professionals with full knowledge of patient history and concerns to assist in their design of an appropriate healthcare program.